

**NEW UNIVERSITY OF LISBON
FACULTY OF MEDICAL SCIENCES**



**“KNOWLEDGE AND PRACTICES OF GENERAL PRACTITIONERS OF
DISTRICT PESHAWAR ABOUT SCHIZOPHRENIA”**

**DR. MUHAMMAD IRFAN
MBBS, MCPS (Psychiatry)**

MASTER THESIS IN INTERNATIONAL MENTAL HEALTH

**SUPERVISOR
PROF. DR. JOSE MIGUEL CALDAS DE ALMEIDA**

Lisbon 2012

ABSTRACT

Schizophrenia with its disabling features has been placed in the top ten of global burden of disease and is associated with long-term decline in functional ability. General Practitioners not only have an important role in treating patients with an established diagnosis of schizophrenia but they can also contribute significantly by identifying people in early stages of psychosis as they are the first hand medical help available and the duration of untreated psychosis is a good indicator of patient's prognosis.

This cross sectional survey, conducted at the clinics of General Practitioners, was designed to assess the knowledge and practices of general practitioners in Peshawar on diagnosis and treatment of schizophrenia. A semi structured questionnaire was used to assess their knowledge and practices regarding schizophrenia. The Knowledge/Practice was then categorized as good or poor based on their responses to the questions of the administered questionnaire.

Overall, the results showed that the knowledge and practices of general practitioners of district Peshawar were poor regarding schizophrenia and may be responsible for delayed diagnosis, inadequate treatment and poor prognosis.

KEY WORDS: Knowledge, Practice, Schizophrenia, General Practitioner.

DEDICATION

*This effort is dedicated to my Family, Friends and
Teachers for their patience and showing great love &
affection for me.*

ACKNOWLEDGMENT

All glory is to Almighty Allah, who gave me good health, opportunity and courage to accomplish this study. I am profoundly indebted to Prof. Dr. Jose Miguel Caldas de Almeida who kindly supervised me throughout my work and critically reviewed the work. His guidance, encouragement and keen interest made the completion of this project possible.

I also wish to thank the supervisor for my clinical studies Prof. Dr. Saeed Farooq who I have always looked up for advice and he has always been kind to guide me in the field of Psychiatry and Research. Also, I am thankful to all my friends for their efforts in searching the appropriate literature and helping me throughout.

I acknowledge my family, for their encouragement and showing special interest in whatever I do.

Dr. Muhammad Irfan

CONTENTS

S. No	Particulars	Page No
	PART-I	
1.	Title	
2.	Abstract	II
3.	Dedication	III
4.	Acknowledgment	IV
5.	Table of contents	V-VII
6.	List of abbreviations	VIII
	PART-II	
	INTRODUCTION	1
	CHAPTER 1: LITERATURE REVIEW	
1.1	General Practice in Pakistan	4
1.2	Introduction to Schizophrenia	6
1.3	Epidemiology of Schizophrenia	7
1.4	Symptoms of Schizophrenia	10
1.4.1	<i>Positive Symptoms</i>	
1.4.2	<i>Negative Symptoms</i>	
1.4.3	<i>Other Symptoms</i>	
1.5	Diagnosis Of Schizophrenia	13

1.6	Aetiology of Schizophrenia	14
1.6.1	<i>Neurodevelopmental and Genetic causes</i>	
1.6.2	<i>Environment and social causes of Schizophrenia</i>	
1.6.3	<i>Dopamine Hypothesis of Schizophrenia</i>	
1.6.4	<i>Structural Brain Changes</i>	
1.7	Course of Schizophrenia	16
1.7.1	<i>Mortality</i>	
1.7.2	<i>Comorbidity</i>	
1.8	Management of Schizophrenia	18
1.8.1	<i>Psychosocial Interventions</i>	
1.8.2	<i>Pharmacological Treatment</i>	
1.9	General Practitioners and Schizophrenia	21
	CHAPTER 2: ORIGINAL STUDY	
2.1	Objectives	23
2.3	Subjects and Methods	
2.3.1	<i>Sample</i>	24
2.3.2	<i>Questionnaire</i>	
2.3.3	<i>Analysis</i>	
2.4	Results	
2.4.1	<i>Knowledge about frequent symptoms of schizophrenia</i>	28
2.4.2	<i>Schizophrenia treatment practices</i>	
2.5	Discussion	32
2.6	Conclusions	40

	CHAPTER 3: TABLES AND FIGURES	
Table 1	Characteristics of the General Practitioners surveyed	41
Table 2	Composite knowledge scores of the General Practitioners	42
Table 3	Correct responses of General Practitioners on multi item knowledge questions	43
Table 4	Composite practice scores of the General Practitioners	44
Table 5	Correct responses of General Practitioners on multi item practice questions	45
Figure 1	Knowledge scores on the frequent symptoms of schizophrenia identified by the Practitioners	46
Figure 2	The distribution of composite knowledge scores of the Practitioners on schizophrenia diagnosis and treatment	47
Figure 3	The distribution of composite practice scores of Practitioners on schizophrenia diagnosis, treatment and referrals	48
	CHAPTER 4: LIST OF REFERENCES	
4.1	References	49
	CHAPTER 5: ANNEXURE	
5.1	Proforma on knowledge and practices of general practitioners of district Peshawar about Schizophrenia	83

LIST OF ABBREVIATIONS

CPSP	College of Physicians and Surgeons Pakistan
DALYs	Disability-Adjusted Life Years
DSM-IV	Diagnostic and Statistical Manual of Mental Disorders
GP	General Practitioner
ICD-10	International Classification of Disease-10
MRI	Magnetic Resonance Imaging
PMDC	Pakistan Medical and Dental Council
SPSS	Statistical Package for Social Sciences
YLDS	Years of Life Lived With Disability

INTRODUCTION

Schizophrenia is a disorder which involves chronic or recurrent psychosis and can lead to long-term decline in functional ability¹. It is placed in the top ten causes of disability in the Global Burden of Disease because it has an early adulthood onset, lifelong course, lack of social acceptability and incapacitating symptoms, which collectively make it one of the most disabling and financially catastrophic disorders². In 1990, the estimated loss in DALY's due to schizophrenia and associated disorders was around 13 million representing almost 1% of burden of the disease from all causes and was ranked 26th in the list². By the year 2020, Schizophrenia is projected to be in 20th position with a DALY's loss of more than 17 million and 1.25% of the overall burden². It is estimated to have a lifetime risk of 0.2 to 0.7%³, with 11/ 100,000 as an annual incidence⁴. The course of illness for an individual patient is difficult to predict. About 10 % of patients recover from an initial episode and do not experience any further impairment but the majority i.e., 55 % has chronic symptoms and the remainders experience an intermittent course⁵. Relapse of psychosis is highly associated with discontinuation (& non compliance) of antipsychotic medication, as well as with substance abuse, psychosocial stressors, and physical illness⁶⁻¹¹.

Schizophrenia affects about 24 million people worldwide with more than 50% of these, not receiving appropriate care and out of 90 % of these, residing in developing countries, Pakistan being one of them¹². Variation in the prevalence of schizophrenia across geographic regions, populations and ethnic groups has been suggested but not confirmed¹³. The exact prevalence of schizophrenia in Pakistan is not known which prevents the making of national level strategies to combat this incapacitating and burdensome illness. Taking in account the global prevalence and the draft of “Assessment of Health Status & Trends in Pakistan”, the estimated prevalence of schizophrenia in Pakistan may be 1-2% in the general population¹⁴ although it varies among rural and urban population of various provinces. In Punjab, it is 2.5% for urban while that in rural is 2%. In Sindh, it is 2% for urban while 1.5% for rural population. In Khyber Pakhtunkhwa, it is estimated to be 2% in urban and 2.5% for rural population. Baluchistan has the lowest with 1% each in urban and rural population¹⁵.

This is a well known fact that Schizophrenia is best treated in specialty clinics but there is a very low number of Psychiatrists and specialty clinics and centers for the diagnosis and treatment of schizophrenia in Pakistan¹⁵. General Practitioners, therefore, have an important role in treating the cases with an established diagnosis

of schizophrenia¹⁶ as well as in identifying people in the early stages of psychosis¹⁷ as they are mostly the first hand medical help available. They come across the diagnosis in a variety of contexts including initial presentation, provision of support to family members, evaluation of concurrent medical illness, management of medication side effects and primary treatment when specialty options are not available and offer an effortless and non-stigmatizing access to health care, to the people¹⁸. Interestingly, there are only a few studies that focus on the knowledge and practice of General practitioners in dealing with patients of schizophrenia around the globe¹⁹⁻²². Therefore, it was decided to conduct a study to assess the knowledge and practices of General Practitioners of district Peshawar about Schizophrenia to highlight the magnitude of the issue and to address the gaps in knowledge and practice in this context and to make plans for improvement, if required.

In Chapter 1, we will review the existing literature on schizophrenia, the role of general practitioners and the situation of general practice in Pakistan. In Chapter 2, we will present the details of the Original study conducted while Chapter 3 focuses on the tables and figures of the Original study. Chapter 4 and 5 describe the references and the proforma used to assess the knowledge and practices of General Practitioners of district Peshawar about Schizophrenia.

CHAPTER 1: LITERATURE REVIEW

1.1 GENERAL PRACTICE IN PAKISTAN

General practitioners (GPs) constitute the majority of health care providers in most parts of the world, treating major bulk of patients and serving as the back bone of any health care system²³. They make up about 85% of all the registered doctors and are responsible for managing approximately 80% of patients in Pakistan²³. However, studies conducted in Pakistan show lack of knowledge and essential expertise in practice of general practitioners' regarding basic health issues²³⁻²⁵. The situation is even worse in rural areas of Pakistan, comprising of 66% population of the country, as specialists and better health facilities are often concentrated in cities creating an imbalance in health service provision²⁶. So, comparatively, there is a poor quality of care in rural areas and therefore higher chance of misdiagnoses or inappropriate referral.

There can be many reasons including the fact that most of the general practitioners do not possess any additional qualification after graduating from medical school and are not re assessed thereafter for their competency²⁷. This creates professional isolation by virtue of staying away from the teaching atmosphere, having none or few opportunities to improve their existing knowledge²⁷.

General practice in Pakistan is comparatively a new and an underdeveloped specialty of medicine with formal training program by the name of Family Medicine initiated by College of Physicians and Surgeons Pakistan (CPSP) in 1992²⁸. Soon after, this training program got discontinued and has restarted just a few years back²⁸. Till date, a very few number of institutions are recognized for the training and the total number of fellows of CPSP in the subject of Family Medicine are just 43²⁹. So all the other general practitioners in Pakistan, unless qualified from abroad with degrees like Members of Royal College of General Practitioners, are just medical school graduates. They need regular program of continuous medical education formulated by Pakistan Medical and Dental Council, the national health regulatory authority, followed by establishing a competency assurance system to ensure the best possible health care delivery to the public³⁰.

Like other developing countries, the number of Psychiatrists for the diagnosis and treatment of schizophrenia is very low¹⁵. Thus, Pakistan is a good case to advocate the importance of the role of general practitioners in this regard^{16, 17} as they can help in treating the psychiatric disorders, which are reported as the third most common reason for consultation in primary care, in an unstigmatized manner^{31, 32}.

1.2 INTRODUCTION TO SCHIZOPHRENIA

Schizophrenia, apart from being in the top ten list of the Global Burden of Disease ^{2, 33}, is also ranked 6th as a cause of disability worldwide, measured by Years of Life lived with Disability³⁴. The word schizophrenia, roughly translated as “splitting of the mind” is derived from the Greek roots schizein “to split”, and phren “mind” ^{35, 36}. A detailed case report by John Haslam and accounts by Phillipe Pinel in 1809, are often considered as the earliest reported cases^{37, 38}.

Schizophrenia is estimated to reduce the life expectancy by approximately 10 years. Schizophrenia, according to the Global Burden of Disease Study, causes a high degree of disability accounting for 1.1% of the total disability-adjusted life years (DALYs) and 2.8% of years lived with disability (YLDs). In the age group 15–44 years, it is the 8th leading cause of DALYs worldwide, according to the World Health Report ³⁹.

Although significant advancement has been attained in the diagnosis, treatment and the disorder’s neurobiological substrates, a comprehensive knowledge of its origins and pathogenic mechanisms is yet to be acquired⁴⁰.

1.3 EPIDEMIOLOGY OF SCHIZOPHRENIA

Psychotic disorders (divided over 12 different diagnostic categories) have a total lifetime prevalence of 3.5%⁴¹. For Schizophrenia, the incidence is relatively low (11 to 15.2 per 100,000)^{4, 42} with wide variation of rise and fall in different populations including those with co morbidities^{43, 44} but the prevalence remains substantially variable across populations, ethnic groups and geographic regions with the consideration of using a different type of prevalence estimate^{45, 46}, tendency to start early in adult life and become chronic⁴⁷. The range of 1-2.5% has also been reported in various areas of Pakistan¹⁵.

At an average, it is accepted that schizophrenia affects 1% of the population, with similar rates across different countries, cultural groups, and sexes⁴⁸.

Schizophrenia typically presents in early adulthood and has a tendency to develop between 16 and 30 years of age, mostly persisting throughout the patient's lifetime⁴⁸. An increase in onset after puberty, which continues throughout adolescence and peaks in the twenties, is reported which then start decreasing from the thirties and tends to tail off in the fifties⁴⁹.

Childhood-onset schizophrenia is defined by an onset of psychotic symptoms before 13 years of age⁵⁰. The illness occurs at a younger age in those with positive family history of schizophrenia⁵¹. Since the

practice of psychiatry in Pakistan is mostly “General Adult” based, a patient of schizophrenia with a childhood or adolescent onset may be too late to present, in terms of prognosis. General Practitioner may have a very important role in diagnosing these cases early when such patients are brought to them with psychotic symptoms attributed to various cultural issues.

With a gender ratio of 1.4:1 male: female, Schizophrenia is more common in men than women^{44, 52}. Studies show that the age of onset of disease is earlier in males than in females by 3 to 5 years, regardless of culture⁵³. Typically, men exhibit symptoms at an earlier age with a worse prognosis. The peak age of onset is in the early twenties, with very few cases occurring after 45 years of age⁵⁴.

The presence of Schizophrenia in African and Caribbean people residing in the United Kingdom, compared to the native white population was reported to be 6 times drawing attention to cultural considerations⁵⁵. The rates are higher in the children of migrants⁵⁶. This may suggest that either the perceived discrimination of the ethnic minority groups or some factor closely related to it is contributing to their increased risk of schizophrenia⁵⁷.

In the “Ethnic Minority Psychiatric Illness Rates in the Community” (EMPIRIC) study, the prevalence of psychosis, among participants of Caribbean, Irish, Bangladeshi, Pakistani and Indian ethnicities, was

three times higher in those experiencing verbal racism, and five times higher in those having a racist physical attack, than in those who did not, supporting the above mentioned social hypothesis⁵⁸.

There is a two way relationship between Schizophrenia and unemployment/ singleness as these increases the risk of developing schizophrenia and subsequently schizophrenia increases the risk of unemployment and singleness⁵⁹. Even in the developed countries, although employment rates may be higher for them but they still are unable to fully support themselves⁶⁰. In Pakistan, psychiatric disorders are more common in population with low socioeconomic status in rural areas and dominantly involve the males⁶¹.

Area of residence also has an important effect on the rates of schizophrenia. Residing in areas of increasing population density increases the hospitalization risk in men with vulnerability for schizophrenia which is expressed as poor cognitive and social abilities^{62, 63}.

There is evidence that schizophrenia is more common in those born in cities, and that the larger the city and the longer a person has lived there, the greater the risk⁴². Schizophrenia is also thought to be related to lower social class⁶⁴. Material deprivation is also likely to influence admission rates for psychosis but social fragmentation has the greatest effect⁶⁵.

1.4 SYMPTOMS OF SCHIZOPHRENIA

In general population, 4.4% report incident psychotic symptoms⁶⁶. There is mostly a prodromal period before the onset of psychosis which is characterized by various mental disturbances like negative symptoms, attenuated and brief transient frank psychotic symptoms, cognitive impairments, and a marked decline is observed in social functioning and quality of life⁶⁷.

Eugen Bleuler described the main symptoms of schizophrenia in 1908 as 4 A's: flattened Affect, Autism, impaired Association of ideas and Ambivalence⁶⁸. Kurt Schneider, a German psychiatrist considered certain symptoms as characteristic of schizophrenia and thus called these as "first rank symptoms"^{69, 70}.

The characteristic symptoms of schizophrenia fall into broad categories of positive and negative symptoms with other associated symptoms of cognitive impairment and affective disturbance³³.

1.4.1 POSITIVE SYMPTOMS

These are synonymous with psychosis. "Positive" refers to the active quality of these symptoms, whose presence is abnormal. Positive symptoms are correlated with first-time diagnosis and hospital admission, but have little predictive value for long-term course. Delusions are the most common symptom, occurring in 65 percent of patients^{71, 72}. Hallucinations and thought disorganization each are

described in about 50 percent of patients ^{73, 74}. A large number of patients experience a combination of delusions, hallucinations, and disorganization⁷¹. Positive symptoms are the most responsive to pharmacological treatment, though they may wax and wane ⁷⁵.

1.4.2 NEGATIVE SYMPTOMS

Negative symptoms represent the diminution or absence of normal characteristics and include flat/ blunted affect and emotion, alogia (poverty of speech), anhedonia (inability to experience pleasure), asociality (lack of desire to form relationships), and avolition (lack of motivation) ^{33, 76}. These deficits may occur months or years before the onset or detection of psychotic symptoms ⁷⁷ and are moderately correlated with functional incapacity, particularly at work^{78, 79}. Evidence suggests that patients suffering from schizophrenia often exhibit a normal or even increased level of emotionality, especially in response to negative events⁸⁰. Contradictory evidence shows that patients with schizophrenia experience both receptive and expressive deficits^{81, 82}. That is, they not only appear blank to others, but see the people around them in the same way.

1.4.3 OTHER SYMPTOMS

Schizophrenia is associated with a wide range of deficits in neurocognitive function including attention, memory, language and executive function^{83, 84}. Significant deficits are present by birth,

followed by moderate decline additionally with the onset of active illness, in most cases⁸⁵⁻⁸⁷. School performance and cognitive testing of individuals who later developed schizophrenia, when reviewed retrospectively, showed a pattern of poor performance^{88, 89}. Cognitive problems are highly correlated with functional impairment of the patient⁹⁰.

The combination of inappropriate, odd and blunted expression is the most frequently observed affective disturbance and is stigmatizing in social settings⁹¹. In early schizophrenia, depressive mood is reported in upto 81% of individuals with first-episode while depression is reported in about 22% of those with first-episode^{92, 93}. Suicide is also reported commonly either at the beginning of treatment or after the resolution of an acute episode or while shifting from hospital to outpatient care⁹⁴. Overall, mood disturbance occurs at about four times the rate seen in the general population⁹⁵.

Few other symptoms worth mentioning include disorganization syndrome (chaotic speech, thought, and behavior)⁹⁶, catatonia (an altered state of motor activity and attention), echopraxia (inappropriately mirroring movements), echolalia (repeating speech in a rigid and stereotypic way), stereotypic and bizarre movements, stuporous appearance or may be internally preoccupation with other psychotic symptoms³³.

1.5 DIAGNOSIS OF SCHIZOPHRENIA

The presentation of schizophrenia varies significantly with a wide range of psychotic manifestations and varying levels of functional incapacity. Schizophrenia is a challenging diagnosis because there are neither pathognomonic features nor confirmatory laboratory or neuropsychological tests.

The most widely used diagnostic criteria for schizophrenia were developed by WHO (International Classification of Diseases, ICD-10) and American Psychiatric Association (Diagnostic and Statistical Manual of Mental Disorders, DSM-IV-TR) with little differences. The ICD-10 criteria emphasize more on Schneiderian First Rank Symptoms suggested by Kurt Schneider^{97, 98}. The ICD 10 includes Paranoid Schizophrenia; Hebephrenic Schizophrenia; Catatonic Schizophrenia; Undifferentiated Schizophrenia; Residual Schizophrenia; Post Schizophrenic Depression; Simple Schizophrenia; under the sub category F 20⁹⁸⁻¹⁰⁶.

1.6 AETIOLOGY OF SCHIZOPHRENIA

The main risk factors are genetic causes, pregnancy and delivery complications, slow neuromotor development, and deviant cognitive and academic performance¹⁰⁷.

1.6.1 NEURODEVELOPMENTAL AND GENETIC CAUSES

Recent studies suggest that abnormalities can be observed years before the onset of positive symptoms¹⁰⁸.

The concordance rate for schizophrenia between monozygotic twins is 50 percent¹⁰⁹. Studies have identified possible gene associations e.g., neuregulin-1 gene, presence of a susceptibility gene (ZNF804A) and increase in gene structural variants¹¹⁰⁻¹¹⁴. It is now widely suspected that there is no single genetic determinant and multiple genetic factors work in combination to create the vulnerability¹¹⁵.

Advanced paternal age is more prevalent in schizophrenia patients and insults to fetal development occurring in first and second trimester and exposure to psychoactive substances, especially cannabis are correlated with an increased risk¹¹⁶⁻¹²².

1.6.2 ENVIRONMENT AND SOCIAL CAUSES

Urbanization, social and racial adversity, family dysfunction and unemployment have all been proposed as risk factors^{48, 63, 123}.

It is suggested that winter or spring birth and prenatal exposure to infections is associated with developing schizophrenia later^{124,125}.

Childhood experiences of abuse or trauma and unsupportive dysfunctional parental relationships have also been noted as risk factors for a diagnosis of schizophrenia^{126, 127}.

There is a two way relationship is proposed to exist between schizophrenia and drugs of abuse or alcohol¹²⁸.

1.6.3 DOPAMINE HYPOTHESIS OF SCHIZOPHRENIA

It has been suggested that positive symptoms are due to hyperactivity of dopaminergic projections from the midbrain while negative and cognitive symptoms are correlated with a decrease in prefrontal activity of dopaminergic pathways^{129, 130}.

1.6.4 STRUCTURAL BRAIN CHANGES

Schizophrenic brains are smaller than normal brains, with ventricular enlargement and thinning of neuritic processes without loss of neuronal bodies¹³¹⁻¹³³. Medial temporal lobes are found to be smaller in the patients of schizophrenia^{134, 135}.

1.7 COURSE OF SCHIZOPHRENIA

Schizophrenia has a quite consistent natural history and longitudinal course. Many patients treated in their first episode of schizophrenia show a good response to treatment and achieve some symptom remission and level of recovery, but recurrent episodes lead to significant neurological deterioration¹³⁶. The role of General practitioner is vital in the early diagnosis of such cases where they sometimes receive the patients with other co morbidities.

Few patients once diagnosed of having schizophrenia, enjoy complete remission. Living in a house hold of 3 or more adults, later age of onset and taking antipsychotics predict complete remission¹³⁷.

The overall rate of recovery during the early years of the illness is low but some patients with first-episode schizophrenia can achieve sustained symptomatic and functional recovery¹³⁸. Patients in remission require markedly less health care resources¹³⁹.

It is also a well known fact that premorbid functioning is associated with better response to treatment, fewer extra pyramidal symptoms and better recovery¹⁴⁰. The duration of untreated psychosis also significantly affects the course of the illness, its symptom severity

and outcome of the illness. The longer the duration of untreated psychosis, the poorer is the prognosis¹⁴¹.

In a nutshell, Schizophrenia is associated with a 20 percent reduction in life expectancy and worse physical health than in the general population^{142, 143}.

1.7.1 MORTALITY

Patients with schizophrenia have a higher mortality and nearly a quarter of deaths resulting from unnatural causes¹⁴⁴. Suicide rates in schizophrenia and other psychotic disorders appear to be 20-fold higher¹⁴⁵. At the same time deliberate self harm or suicide attempt is also thought to be a predictor of relapse¹⁴⁶.

1.7.2 COMORBIDITY

The most common comorbid condition is substance abuse. As many as 80 percent of patients of schizophrenia abuse alcohol, illicit drugs, or prescription medications¹⁴⁷⁻¹⁴⁹.

Patients of schizophrenia with comorbid cannabis abuse have more positive symptoms and show more violent behaviour¹⁵⁰.

Increased rates of chronic medical conditions like coronary artery disease, chronic obstructive pulmonary conditions, hepatitis, hypothyroidism, diabetes mellitus, fluid and electrolyte disorders are observed in patients with schizophrenia¹⁵¹.

1.8 MANAGEMENT OF SCHIZOPHRENIA

The introduction of chlorpromazine in the 1950s, changed the complete scenario of management of schizophrenia as previously, the care of patients suffering from schizophrenia was limited to various rehabilitative, psychotherapeutic and custodial interventions. The advent of relatively safe and effective pharmacologic treatments paved the path for marked improvement in symptoms and functioning and made it possible for the majority of the patients to live in community settings¹⁵². Effective treatment interventions using a combination of best possible pharmacotherapy and targeted psychosocial treatments are elevating expectations about the prospects of functional recovery in patients with schizophrenia¹⁵³.

1.8.1 PSYCHOSOCIAL INTERVENTIONS

There are various guidelines for the treatment of schizophrenia developed across the world. These recommend nearly similar pharmacotherapy but have variations in the type of psychosocial interventions which describes their importance^{154, 155}.

Psychological interventions for schizophrenia include cognitive behavioral therapy for symptoms, cognitive remediation for neurocognitive deficits, motivational interventions for substance misuse and for non-adherence to medication and family

interventions¹⁵⁶. Social skills' training is planned to deal with the deficits in patient's communication and social interactions¹⁵⁷. A recent study, reported in Pakistan showed promising preliminary results with the use of CBT in patients with psychosis¹⁵⁸.

Vocational rehabilitation is useful at getting patients into the place of work, although they may rarely work in a long-term competitive employment¹⁵⁹.

1.8.2 PHARMACOLOGICAL TREATMENT

There are three basic classes of medications (typical, atypical and dopamine partial agonist antipsychotics) which act principally on dopamine systems¹⁶⁰. The typical antipsychotic agents have been associated with relatively high incidence of adverse effects ranging from acute dystonia to akathisia and akinesia to tardive dyskinesia and neuroleptic malignant syndrome¹⁶¹. Atypical agents with greater affinities for serotonin and norepinephrine¹⁶² lead the patients to receive significantly less prescriptions for anticholinergics¹⁶³. Clozapine, dopamine partial agonist, remains the treatment of choice for refractory schizophrenia, although it is well known to cause blood dyscrasias and other serious adverse effects such as seizures, intestinal obstruction, myocarditis, thromboembolism and cardiomyopathy¹⁶⁴.

Obesity, type 2 diabetes mellitus, and hyperlipidemia (the metabolic syndrome) occur both with schizophrenia and with antipsychotic medications¹⁶⁵⁻¹⁶⁸. Cardiac risk associated with schizophrenia and with antipsychotic drugs has recently been fully appreciated¹⁶⁹⁻¹⁷¹. Antipsychotic medications vary in their propensity to cause QT_c prolongation but a particular concern is with intravenous use of haloperidol, which has led to fatality and a 2007 FDA alert¹⁷².

In Pakistan, drug non-compliance has been a major hindrance in the effective management of schizophrenia. Nearly 74% of the patients have a relapse of the illness and need frequent readmissions resulting from the non-compliance¹⁷³. Non-affordability of drugs, unawareness of the benefits of treatment, physical side effects and unfriendly attitude of the doctors¹⁷⁴ are the commonest reasons for non-compliance.

1.9 GENERAL PRACTITIONERS AND SCHIZOPHRENIA

In Low and Middle Income countries, most people with schizophrenia probably receive little or no formal care. One manifestation of this is very long duration of untreated psychosis in the first episode, i.e., 132 weeks¹⁷⁴. This poses a major public health problem considering that around 41.7 million people with schizophrenia may need care in these countries¹⁷⁵. The general practitioners, therefore, have to play a crucial role in early diagnosis and management, as they come across these cases in various contexts. To fulfill the role, the general practitioners need diagnostic knowledge, low-threshold, easily accessible specialized services to which they can refer these patients.

The movement, and then the subsequent policies in favour of closing the mental hospitals and transferring these to community services have been campaigned for quite some time¹⁷⁶ and are being actively pursued recently¹⁷⁷. This may also increase the role of general practitioners being the physicians readily available in the community. Guidelines for schizophrenia management in general practice emphasize on the early diagnosis of psychosis, minimizing delays in treatment initiation, keeping an eye on the patient's condition and treatment adherence, and prompt intervention at times of relapse or psychosocial crises¹⁷⁸⁻¹⁸⁴. The early diagnosis of psychosis, thus,

becomes very important as it is the central part of “mental health literacy” which is mandatory for appropriate help-seeking¹⁸⁵, and the general practitioners can play a pivotal role in this regard.

General practitioners may be less confident in the clinical skills than a psychiatrist regarding schizophrenia but can see themselves complementing the psychiatrist with an active and useful role¹⁸⁶. The level of confidence can be improved by including mental health in the undergraduate curriculum and by providing pre service training to the general practitioners. Once they feel confident about the knowledge, the universal screening should be emphasized to improve detection rates of mental disorders including schizophrenia for which brief mental health screening questionnaires might help¹⁸⁷.

Unfortunately, there are only a few studies that focus on the knowledge and practice of General practitioners about schizophrenia, around the globe¹⁹⁻²². The authors are not aware of studies from other developing countries which address this issue except a study in a semi urban small border district in Pakistan with a smaller sample size using the same proforma²². It was therefore decided to conduct a study to assess the knowledge and practices of General Practitioners about schizophrenia in a major urban center, Peshawar, to identify the gaps in knowledge and practice in service delivery, and chalk out plans to improve it, if needed.

CHAPTER 2: ORIGINAL STUDY

2.1 OBJECTIVES

To assess the knowledge and practices of general practitioners in district Peshawar on diagnosis and treatment of schizophrenia.

To estimate the frequency of patients of schizophrenia, seen in general practice in district Peshawar.

2.2 SUBJECTS AND METHODS

2.2.1 Sample

This cross sectional survey was conducted at the clinics of General Practitioners in Peshawar from August 2009 to December 2011. A list of General Practitioners of Peshawar, enrolled with provincial Health Regulation Authority was obtained and all the 135 listed General Practitioners were contacted for the purpose of the survey. Unlike the countries such as UK, in Pakistan the GPs are not defined by their registration or after a specified period of training. For the purpose of this study a General Practitioner was defined as, "A licensed medical graduate registered with Health Regulatory Authority who gives personal, primary and continuing care to individuals, families and a practice population irrespective of age, sex and illness"¹⁸⁸.

All the enlisted (n=135) were approached to give consent to participate in the study. Out of 135, 114 consented to participate in the study and were included through purposive, non-probability sampling. The study was given ethical approval by the institutional review and ethical board, Postgraduate Medical Institute, Lady Reading Hospital Peshawar.

Twenty One General Practitioners did not consent to participate in the study.

It is worth mentioning that most of the general practices in Peshawar are single doctor based practices providing service for fee. The GPs only provide curative services and patients meet the treatment costs through out of pocket expenses.

2.2.2 Questionnaire

The participating General Practitioners were then requested to answer a semi structured questionnaire (Appendix A) consisting of three parts namely, General Information, Knowledge and Practices related to Schizophrenia. This questionnaire has been used by Simon AE et al in their seminal study and has been validated for use with GPs¹⁹. As this questionnaire was developed in Switzerland where the health system and the practice of GP differs markedly from that in Pakistan, the questionnaire was modified to reflect the practice and health services in Pakistan. However, we retained the major domains, questions and content as developed by Simon AE et al¹⁹. The 17 item questionnaire consisted of 5 demographic items and 12 (partly multi-item) questions that assessed: Knowledge (symptoms of schizophrenia; early warning signs of schizophrenia; treatment; and management of schizophrenia patients) and Practice (methods used to confirm diagnosis; referrals used; and medications prescribed to schizophrenia patients) and continuing medical education of the Practitioner.

Knowledge was defined as the ability of the general practitioner to correctly identify the symptoms of schizophrenia and its related treatments and appropriate referrals. Similarly practice was defined as the ability of the general practitioner to correctly indicate their practice in diagnosing, treating and referring schizophrenia patients appropriately. The level of knowledge on schizophrenia diagnosis and treatment was assessed with two multi-item questions (K1 and K3) and five other questions (K2, K4-K7) having a maximum score of 19. A discriminating index was developed using the composite scores obtained by the general practitioners' responses to all knowledge questions. Good knowledge was defined as a composite knowledge response score of $\geq 60\%$ and Poor Knowledge was defined with a composite knowledge score of $< 60\%$. The practical experience of general practitioners with schizophrenia diagnosis, treatment and referrals was assessed with four multi-item questions (P1 – P4) having a maximum score of 22. A similar discriminating index using the composite scores obtained by the general practitioners' responses to all the Practice questions was developed. Adequate practice was defined as a composite practice response score of $\geq 60\%$ and Inadequate practice was defined with a composite practice response score of $< 60\%$.

The questionnaires were distributed in the clinics of General Practitioners along with a short briefing about the questionnaire. The questionnaires were then left at their clinics and a collection time was decided with them. The filled questionnaires were collected by re-visiting their clinics in person. However, not all the GPs provided the filled questionnaire on the decided date and there have been instances where the collection of the filled questionnaire needed many visits.

The scoring scheme based on the correct answers was developed in consultation with the data analyst. All the questions and the items of the multi-item questions regarding the knowledge and practice were marked as either correct (score=1) or incorrect (score=0).

2.2.3 Analysis

Data collected through semi structured questionnaire was analyzed using Epi Info Statistical software. The demographic details were calculated using percentages, and proportions were calculated using the discriminating index of composite scores for Knowledge and Practices of the Practitioners regarding schizophrenia.

2.3 RESULTS

There were 135 General Practitioners invited to participate in this study, in that 114 (84.44%) consented to participate and 21 (15.56%) declined to participate in the study. Majority of GPs were male (n=111, 97%).

About 13% (n=15) of the General Practitioners in this study treat more than 10 schizophrenia patients annually and a majority of them see few to none of the schizophrenia patients in Peshawar. None of them did possess any specialty training in mental health and very few (n=7, 6.1%) General Practitioners received continuing medical education regarding schizophrenia (Table 1).

The 21 General Practitioners who refused to participate gave reasons for their non-participation including “I have a busy schedule” 13 (61.9%); “I don’t see patients with schizophrenia” 6 (28.6%); and “I don’t want to fill the questionnaire” 2 (9.5%).

2.3.1 KNOWLEDGE ABOUT FREQUENT SYMPTOMS OF SCHIZOPHRENIA

The level of Schizophrenia Diagnostic knowledge of Practitioners was assessed through multiple questions. A composite score on the responses to all the seven knowledge questions (K1-K7) was estimated with the response scores ranging from 0 to 16 (Figure 2). Good knowledge was identified among 12.2% (n=14) of the

surveyed general practitioners with a composite score of ≥ 12 . About 6% (n=7) of the general practitioners had no knowledge at all about the diagnosis and treatment of schizophrenia receiving a composite knowledge score of 0 and the remaining 81.6% (n=93) of general practitioners had very poor knowledge (Table 2).

The frequent symptoms of schizophrenia were identified correctly ($\geq 60\%$ correct response) by 31.5% (n=36) of the Practitioners surveyed, with a response score of ≥ 6 . About 60% (n=69) had poor knowledge about the frequent symptoms of schizophrenia while 7.9% (n=9) appeared to have no knowledge about the disease (Figure 1). A large number of general practitioners (n=93, 81.6%) considered hallucinations and delusions to be the most frequent symptoms of Schizophrenia (Table 3).

Eighty (70.2%) general practitioners considered pharmacotherapy to be ideal for a patient with a suspected first schizophrenic episode (Table 3).

Seventy one (62.3%) general practitioners, each, considered that the first episode of schizophrenia is preceded by early warning signs and estimated the relapse risk of untreated patients during the first year after a first schizophrenic episode to be more than 60 percent. Only 15 (13.2%) considered giving antipsychotic medication for 12-24 month after a first schizophrenic episode and 16 (14.0%) considered

giving it for 3-5 years for maintenance therapy in patients with multiple episodes of illness after the remission of an episode.

Only 6 (5.3%) general practitioners could name two side effects and 25 (21.9%) reported one side effect while 83 (72.8%) could not report any side effect of antipsychotic medication. Extra Pyramidal Symptoms were the most relevant side effect reported by 27 (23.7%) general practitioners followed by weight gain reported by 4 (3.5%).

2.3.2 SCHIZOPHRENIA TREATMENT PRACTICES

A composite score on the responses to all the four practice questions (P1-P4) was estimated with the response scores ranging from 0 to 18 (Figure 3). Adequate practice was identified among 28.1% (n=32) of the surveyed general practitioners with a composite practice score of ≥ 13 . About 8.8% (n=10) of the general practitioners did not conform at all to the correct practice guidelines for diagnosis and treatment of schizophrenia receiving a composite practice score of 0 and the remaining 63.1% (n=72) of general practitioners had inadequate practice methods employed in their clinics with a score of less than 13 (Table 4).

Regarding the breakdown of the practice questions about schizophrenia, a sizable majority of general practitioners (n=87, 76.3%) relied on personal history and observation over several

months for the confirmation of the diagnosis of schizophrenia. On the subject of the place of treatment, 68 (59.6%) general practitioners were of the opinion of treating patients of schizophrenia exclusively in their clinic while 45 (39.5%) considered referral to a specialist/psychiatric out-patient department and complete handover for treatment. Only 3 (2.6%) general practitioners could name four medications with their average doses while 66 (57.9%) were not able to name any medication for the treatment of schizophrenia. Forty three (37.7%) general practitioners correctly reported that prognosis of a treated patient after a first episode of schizophrenia is favourable after single episode with a possibility of maintenance of performance level (Table 5).

2.4 DISCUSSION

World Health Organization has repeatedly reiterated that most psychiatric disorders in developing countries should be treated in primary care in developing countries. This is necessitated not only by the very inadequate specialist services but this also helps to reduce stigma and institutionalization. This is also consistent with the fact which emphasize that “early detection of psychosis, minimizing delays in obtaining treatment, monitoring the patient’s condition and adherence to treatment, and prompt intervention at times of relapse or psychosocial crises”^{178, 179, 187, 189-192}. Keeping this in consideration, the knowledge and practices of general practitioners appear to have an important role in managing patients with schizophrenia. This is even more appropriate when we consider the unavailability of community services in our set up¹⁹³.

The limitations of this survey should be kept in mind. The assessment of knowledge and practice was based on a cross sectional view of the GPs responses to a questionnaire. The sample in this study may not be representative of the GPs. The results of our study may not be generalisable to other settings as the training and role of general practitioners may vary in the international healthcare system¹⁹. However, we had a good response rate and were able to contact most of the GPs working in Peshawar District which was

even better than a similar study conducted in a small semi urban border district of Pakistan²².

Majority of respondents were male GPs .This perhaps reflects limited number of female doctors who are mostly limited to providing obstetric services.

International multicentre survey of general practitioners indicates that the mean number of patients with chronic schizophrenia seen by general practitioners is similar across several countries and healthcare systems¹⁹⁴. Although, little is known about general practitioners' experiences in treating schizophrenia, most of them are currently treating patients with schizophrenia in a small number (on average about 3 patients) which is similar to our finding where only 13.2% (n=15) treated more than 10 patients annually as compared to 53.5 % (n=61) general practitioners who didn't treat any diagnosed case of schizophrenia in a year making almost two patients on average^{22, 186, 195}.

It is reported internationally that 40-50% of general practitioners screen patients for mental health issues routinely but depression is often not identified¹⁹³. When a much common condition like depression can be missed often, schizophrenia has all the chances to be missed which was obvious from our findings where 40.4% (n=46) didn't diagnose any new case in a year's time.

In our study, only 6.1% General Practitioners received continuing medical education, an essential ingredient to maintain competence, regarding schizophrenia which, however, is slightly better than none reported in the study by Akhtar et al, but far less than the 30.8% of GPs receiving recent medical education about mental health problems as reported in another study from Pakistan^{22, 196}.

Regarding the knowledge, only 12.3% general practitioners in our sample had adequate knowledge about schizophrenia which visibly speaks about the current state of the affair in Pakistan and is in contrast with the findings of a French study where GPs had a fair theoretical knowledge of schizophrenia symptoms¹⁹⁷.

As compared to the figures reported by the Swiss and a local study (62% and 64% respectively), Hallucinations and delusions were considered as the most frequently experienced symptoms of schizophrenia by 81.6% of the general practitioners in our sample^{19, 22}. However, Bizarre behavior which was considered by 56% in Swiss study was considered by only 45.2% GPs of our sample¹⁹.

General practitioners in our sample (62.3%) considering that the first episode of schizophrenia is preceded by early warning signs were much less than the Swiss study (90%) but comparable to another local study (69%)^{19, 22}. This is an alarming situation as it shows the

unpreparedness of encountering schizophrenia by general practitioners.

Pharmacotherapy, alone or in possible combination with other types of therapy, was considered ideal treatment by 88% of GPs in the Swiss sample and 60% in Akhtar et al sample ^{19, 22}. Our results were in between the two, where 70.2% GPs considered pharmacotherapy to be ideal.

Only 13.2% and 14% GPs of our sample were able to answer correctly the duration of antipsychotic medication to be maintained for more than 12 months after first schizophrenic episode, and for at least 3 years in patients with multiple episodes of illness after remission¹⁹⁸. Figures of 12.5% and 39% respectively were reported for the same in a similar study ²². This is distressing to know as it suggest that even those patients who are diagnosed as cases of schizophrenia, may receive incomplete treatment regime leading to more chances of relapse.

Seventy one percent GPs in our sample correctly estimated the relapse risk of untreated patients during the first year after a first schizophrenic episode which was more than those correctly reported by GPs in other studies ^{19, 22}.

Regarding their practices, the answers given to the questions in the questionnaire may not assess their actual practice, but should be

considered more of a proxy measure¹⁹⁹. Only 20.2% (n=23) of the general practitioners had adequate practice in management of schizophrenia. This percentage was much less than the percentage of any practice question answered by General Practitioners in a similar study conducted in Switzerland¹⁹.

In our sample, 76.3% GPs, each, relied on personal history and observation over several months for the confirmation of the diagnosis of schizophrenia while the GPs in Swiss study showed more reliance on Information from significant others (65%) and Family history (63%) respectively ¹⁹. The sample in the study by Akhtar et al relied more on personal history (87.5%) and family history (70%) respectively ²². All of these serve as important diagnostic predictors of schizophrenia²⁰⁰.

As compared to a figure of 6.9% referrals of all the patients to specialists, 39.5% GPs in our sample showed the same practice which though is much bigger than that reported in the Swiss study, is far less than the 60% reported in another study ^{19, 22}. However 61.4% of the general practitioners were of the opinion of collaborating with the specialists which is comparable to 77.2% in the Swiss survey which is an encouraging finding ¹⁹.

Antipsychotics (one or more) were mentioned as commonly used drug for schizophrenia in their practice by 42.1% of GPs which is far

less than the figures reported in other studies (80% and 98.5% respectively) ^{19, 22}. A study from Pakistan reported that the problem in the rational use of psychotropic medication is due to the gaps in GPs' knowledge about the management of mental disorders¹⁹⁶. This is an area of great concern as it poses serious questions not only on the practice but on the basic medical knowledge of these GPs.

The results of the study are not surprising in view of the fact that teaching and training in Psychiatry is still much limited in undergraduate curriculum in most medical schools in Pakistan. There is little or no training in psychiatry for family physicians and continuing medical education hardly exists. This may well be contributing to a long Duration of Untreated psychosis in the First Episode which is found to be more than two years in developing countries.

Although there are studies on evaluation of the training programmes for GPs^{18, 201} but we believe this is a unique study assessing the knowledge and practice of doctors working in general practice about schizophrenia in a major urban area of a developing country setting. It appears that many educational programmes for GPs have been devised without assessing the prior knowledge, practice and training needs of the GPs. These findings have serious implications for

training the general practitioners in diagnosing and treating the Schizophrenia.

As the overall prevalence of schizophrenia is low, it can be argued that it will be insufficient to improve knowledge and practices of general practitioners only through educational programmes. In addition, synchronized provision of specialized services and assessment facilities for general practitioners may prove to be more important¹⁹. From the general practitioners' point of view, communication between the specialist and primary care services has been poor and problematic with unclear individual roles and responsibilities while treating patients with schizophrenia and this may appear to be the biggest hindrance in service provision^{17, 186}. An integrated system of care should be developed with clearly defined role of GP in the chain of care, which has been a pillar of Mental Health Plan for Pakistan^{202, 203}. Such a system would significantly contribute to the quality of care provided to the patients with Schizophrenia²⁰¹.

Since it has already been established that rapid social integration can be achieved if the follow up of patients with schizophrenia is arranged in general practice, the role of general practitioner for dealing with somatic symptoms of the illness and renewing/changing their anti psychotic medication can't be

emphasized any more^{204, 205}. This will definitely lead to a much needed collaborative care approach^{204, 206-208}.

The use of universal screening by all general practitioners is needed to improve detection rates for which brief mental health screening questionnaires might be very useful and consideration should be given to the development of standardized methods, suggested by GPs during the data collection meetings with them^{17, 193, 209, 210}.

2.5 CONCLUSION

The findings of this study suggest that regarding schizophrenia, the knowledge and practices of general practitioners in a major urban centre in Pakistan are poor and may be one of the reasons responsible for delayed diagnosis and perhaps inadequate treatment. The training and education for general practitioners in the diagnosis and treatment of schizophrenia needs to be improved significantly through more refined, result oriented mental health education. At the same time, communication between mental health services and general practitioners need particular improvement which will not only lead to better access to services but also to knowledge sharing with GPs.

CHAPTER 3: TABLES AND FIGURES

Table 1: Characteristics of the General Practitioners surveyed (n=114)

Characteristics	Number (%)
Gender:	
Male	111 (97.4)
Female	3 (2.6)
Number of schizophrenia patients treated annually:	
None	61 (53.5)
1 – 2 patients	26 (22.8)
3 – 5 patients	8 (7.0)
6 – 9 patients	4 (3.5)
More than 10 patients	15 (13.2)
Time taken to treat a patient:	
Less than 10 minutes	22 (19.3)
10 – 20 minutes	34 (29.8)
21 – 30 minutes	18 (15.8)
More than 30 minutes	6 (5.3)
No response	34 (29.8)
Number of suspected cases of schizophrenia seen annually:	
None	46 (40.4)
1 – 2 patients	38 (33.3)
3 – 5 patients	3 (2.6)
More than 5 patients	27 (23.7)
Continuing Medical Education attended:	
Yes	7 (6.1)
No	107 (93.9)

Table 2: Composite Knowledge scores of the General Practitioners (n=114)

Composite Knowledge Scores	Number (%)
No Knowledge = 0 Score	7 (6.1)
Poor Knowledge = Scores < 12	93 (81.6)
Good Knowledge = Scores \geq12	14 (12.3)

Table 3: Correct responses of General Practitioners on multi item knowledge questions

Multi Item Knowledge Questions		Number (%)
Most frequent symptoms of Schizophrenia		
	Hallucinations/delusions	93 (81.6%)
	Social withdrawal	34 (29.8%)
	Psychosomatic complaints	64 (56.1%)
	Suicidality	68 (59.6%)
	Depression/anxiety	55 (48.2%)
	Bizarre behaviour	52 (45.6%)
	Drug misuse	74 (64.9%)
	Conflicts with parents/teachers/employee	39 (34.2%)
	Functional decline (school/work)	34 (29.8%)
Ideal treatment for a patient suspected of first episode schizophrenia		
	Psychotherapy	17 (14.9%)
	Pharmacotherapy	80 (70.2%)
	Family Therapy	60 (52.6%)
	Observe and Wait only	90 (78.9%)

Table 4: Composite Practice scores of the General Practitioners (n=114)

Composite Practice Scores	Number (%)
No Practice = 0 Score	10 (8.8)
Inadequate Practice = Scores < 13	72 (63.1)
Adequate Practice = Scores \geq 13	32 (28.1)

Table 5: Correct responses of General Practitioners on multi item practice questions

Multi Item Practice Questions	Number (%)
General reliance on confirmation of diagnosis:	
Personal history	87 (76.3)
Family history	75 (65.8)
Information from significant others(teacher/employer)	42 (36.8)
Observation over several days and weeks	78 (68.4)
Observation over several months	87 (76.3)
Neurological assessment	74 (64.9)
Neuropsychological assessment	66 (57.9)
Other examinations (radiographic, electrophysiological)	75 (65.8)
Laboratory tests	77 (67.5)
Urine testing for drug abuse	76 (66.7)
Consultation with/referral to a specialist	39 (34.2)
Treatment alone or in collaboration with other specialists or institutions:	
Treatment exclusively in my clinic	68 (59.6)
Occasional/regular consultation with a specialist to reassess/ advise	33 (28.9)
Referral to a specialist for initial diagnosis and to establish the medication regimen, continuation of treatment in my clinic	37 (32.5)
Referral to a specialist/psychiatric out-patient department and complete handover for treatment	45 (39.5)
Commonly used antipsychotic medications and their doses:	
None	66 (57.9)
One	21 (18.4)
Two	19 (16.7)
Three	5 (4.4)
Four	3 (2.6)
Prognosis of a treated patient after a first episode of schizophrenia:	
Favourable; Single episode with maintenance of performance level is possible	43 (37.7)
Mostly several episodes with possible maintenance of performance level	49 (43.0)
Mostly several episodes with progressive decline of performance level and severe course of illness	31 (27.2)

Figure 1: Knowledge scores on the frequent symptoms of schizophrenia identified by the Practitioners




Figure 2: The distribution of composite knowledge scores of the Practitioners on schizophrenia diagnosis and treatment





Figure 3: The distribution of composite practice scores of Practitioners on schizophrenia diagnosis, treatment and referrals



CHAPTER 4: LIST OF REFERENCES

4.1 REFERENCES

1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 4th ed. Washington DC: American Psychiatry Association; 1994. p.285.
2. Murray CJL, Lopez AD. The Global Burden of Disease. Cambridge: Harvard University Press; 1996. p.21.
3. Kendler KS, Gallagher TJ, Abelson JM, Kessler RC. Lifetime prevalence, demographic risk factors, and diagnostic validity of nonaffective psychosis as assessed in a US community sample. The National Comorbidity Survey. Arch Gen Psychiatry 1996;53:1022-31.
4. Goldner EM, Hsu L, Waraich P, Somers JM. Prevalence and incidence studies of schizophrenic disorders: a systematic review of the literature. Can J Psychiatry 2002;47:833-43.
5. Robinson D, Woerner MG, Alvir JM, Bilder R, Goldman R, Geisler S et al. Predictors of relapse following response from a first episode of schizophrenia or schizoaffective disorder. Arch Gen Psychiatry 1999;56:241-7.

6. Gilbert PL, Harris MJ, McAdams LA, Jeste DV. Neuroleptic withdrawal in schizophrenic patients. A review of the literature. *Arch Gen Psychiatry* 1995;52:173-88.
7. Schooler NR. Relapse prevention and recovery in the treatment of schizophrenia. *J Clin Psychiatry* 2006;67:19-23.
8. Almerie M, Alkhateeb H, Essali A, Matar HE, Rezk E. Cessation of medication for people with schizophrenia already stable on chlorpromazine. *Cochrane Database Syst Rev* 2007; CD006329.
9. Hunt GE, Bergen J, Bashir M. Medication compliance and comorbid substance abuse in schizophrenia: impact on community survival 4 years after a relapse. *Schizophr Res* 2002;54:253-64.
10. Ayuso-Gutierrez JL, del Rio Vega JM. Factors influencing relapse in the long-term course of schizophrenia. *Schizophr Res* 1997;28:199-206.
11. Nuechterlein KH, Dawson ME, Gitlin M, Ventura J, Goldstein MJ, Snyder KS, et al. Developmental Processes in Schizophrenic Disorders: longitudinal studies of vulnerability and stress. *Schizophr Bull* 1992;18:387-425.

12. World Health Organization: Schizophrenia. [Online] 2012 [Cited on March 12, 2012]. Available from URL: http://www.who.int/mental_health/management/schizophrenia/en/.
13. Torrey EF. Prevalence studies in schizophrenia. *Br J Psychiatry* 1987;150:598-608.
14. Department of Community Health Sciences, Aga Khan University: Assessment of Health Status & Trends in Pakistan [Online] 2001 [Cited on April 15, 2012]. Available from URL: <http://www.aku.edu/SiteCollectionImages/AKU/zzchs/pdf/HealthSituationandTrend-Pakistan,2001.pdf>.
15. Gadit AAM. Psychiatry in Pakistan: 1947-2006: A new balance sheet. *J Pak Med Assoc* 2007;57:453-63.
16. Lang FH, Johnstone EC, Murray GD. Service provision for people with schizophrenia. II. Role of the general practitioner. *Br J Psychiatry* 1997;171:165-8.
17. Skeate A, Jackson C, Birchwood M, Jones C. Duration of untreated psychosis and pathways to care in first-episode psychosis: investigation of help-seeking behaviour in primary care. *Br J Psychiatry* 2002;181:S73-7.

18. Oud MJT, Schuling J, Slooff CJ, de Jong BM. How do General Practitioners experience providing care for their psychotic patients? BMC Family Practice 2007; 8:37-43.
19. Simon AE, Lauber C, Ludewig K, Braun-Scharm H, Umbricht DS. General practitioners and schizophrenia: results from a Swiss survey. Br J Psychiatry 2005;187:274-81.
20. Farooq S. Never treated schizophrenia in developing countries. J Pak Psych Soc 2008;5:53-4.
21. Heyrman J, Spreeuwenbergh C. Vocational training in general practice. Leuven: Katholieke Universiteit Leuven; 1987.
22. Akhtar J, Haque M, Idress M, Awan NR, Irfan M. Knowledge and practice of general practitioners about schizophrenia. J Postgrad Med Inst 2010;24:318-22.
23. Marsh D, Hashim R, Hassany F, Hussain N, Iqbal Z, Irfanullah A, et al. Front-line management of pulmonary tuberculosis: an analysis of tuberculosis and treatment practices in urban Sindh, Pakistan. Tuber Lung Dis 1996;77:86-92.
24. Shera AS, Jawad F, Basit A. Diabetes related knowledge, attitude and practices of family physicians in Pakistan. J Pak Med Assoc 2002; 52:465-70.

25. Hussain SF, Zahid S, Khan JA, Haqqee R. Asthma management by general practitioners in Pakistan. *Int J Tuberc Lung Dis* 2004; 8:414-7.
26. Farooq U, Ghaffar A, Narru IA, Khan D, Irshad R. Doctors perception about staying in or leaving rural health facilities in District Abbottabad. *J Ayub Med Coll Abbottabad* 2004;16:64-9.
27. Goulet F, Jacques A, Gagnon R. An innovative approach to remedial continuing medical education, 1992-2002. *Acad Med* 2005;80:533-40.
28. Family Medicine Residency Program: The Aga Khan University. [Online] 2012. [cited 20 January 2012]. Available from <http://www.aku.edu/collegesschoolsandinstitutes/medicine/pakistan/programmes/postgraduate/Pages/residencyfamilymedicine.aspx> URL:
29. List of Good Standing Fellows of CPSP: College of Physicians and Surgeons Pakistan. [Online] 2012 [cited 16 October 2012]. Available from URL: <http://cpsp.edu.pk/index.php?code=b25saW5lX2RpcmVjdG9yeXxmZWxsb3dzX2RldGFpbC5waHB8MHwwfDA=>

30. Rauf MA, Saeed AB. Competency assurance of general practitioners--role of regulatory authority. J Pak Med Assoc 2007;57:573-4.
31. Shah A. The burden of psychiatric disorders in primary care. Int Rev Psychiatry 2001;4:243-50.
32. Akhtar A, Farooq S, Mukhtarul Haq M, Khan SA. The perception and attitude of primary care physicians about psychiatric disorders and their management. J Postgrad Med Inst 2005;19:292-6.
33. Jibson MD, Glick ID, Tandon R. Schizophrenia and Other Psychotic Disorders. Focus 2004; 2:17-30.
34. Farooq S. Free pharmacological treatment for schizophrenia in developing countries- Case for Public Health Intervention. J Pak Psych Soc 2006;3:1-2.
35. Kuhn R, Cahn CH. Eugen Bleuler's concepts of psychopathology. Hist Psychiatry 2004;15:361–6.
36. Schizophrenia. [Online]. 2009. [cited 2009 December 12]. Available from URL:<http://en.wikipedia.org/wiki/Schizophrenia>.
37. Heinrichs RW. Historical origins of schizophrenia: two early madmen and their illness. J Hist Behav Sci 2003;39:349–63.

38. James Tilly Matthews. [Online]. 2009. [cited 2009 February 22]. Available from URL:http://en.wikipedia.org/wiki/James_Tilly_Matthews.
39. Rossler W, Salizeb HJ, Osc JV, Riecher-Rosslerd A. Size of burden of schizophrenia and psychotic disorders. Eur Neuropsychopharmacol 2005;15:399-409.
40. Tsuang MT, Stone WS, Faraone SV. Toward Reformulating the Diagnosis of Schizophrenia. Am J Psychiatry 2000;157:1041-50.
41. Perala J, Suvisaari J, Saarni SI, Kuoppasalmi K, Isometsa E, Pirkola S, et al. Lifetime prevalence of psychotic and bipolar I disorders in a general population. Arch Gen Psychiatry 2007;64:19 -28.
42. McGrath JJ. Variations in the incidence of schizophrenia: Data versus dogma. Schizophr Bull 2006;32:195-7.
43. Boydell J, Van Os J, Caspi A, Kennedy N, Giouroukou E, Fearon P et al. Trends in cannabis use prior to first presentation with schizophrenia, in South-East London between 1965 and 1999. Psychol Med 2006;36:1441-6.
44. McGrath JJ, Saha S, Welham J, El-Saadi O, MacCauley C, Chant DC. A systematic review of the incidence of schizophrenia: The distribution of rate items and the influence

of methodology, urbanicity, sex and migrant status. *Schizophr Res* 2004;67:65-6.

45. Arehart-Treichel J. Global Schizophrenia Prevalence May Have Been Overstated. *Psychiatr News* 2005;40:9-54.
46. Torrey EF. Prevalence studies in schizophrenia. *Br J Psychiatry* 1987;150:598-608.
47. Saha S, Chant D, Welham J, McGrath J. A systematic review of the prevalence of schizophrenia. *PloS Med* 2005;2:413-33.
48. Mueser KT, McGurk SR. Schizophrenia. *Lancet* 2004;363:2063-72.
49. Kirkbride JB, Fearon P, Morgan C, Dazzan P, Morgan K, Tarrant J, et al. Heterogeneity in Incidence Rates of Schizophrenia and Other Psychotic Syndromes. Findings From the 3-Center AEsOP Study. *Arch Gen Psychiatry* 2006;63:250-8.
50. Shaw P, Sporn A, Gogtay GP, Greenstein D, Gochman P, Tossell JW, et al. Childhood-Onset Schizophrenia. *Arch Gen Psychiatry* 2006;63:721-30.
51. Naqvi H, Khan MM, Faizi A. Gender differences in age at onset of Schizophrenia. *J Coll Physicians Surg Pak* 2005;15:345-8.

52. Picchioni MM, Murray RM. Schizophrenia. *BMJ* 2007;335:91-5.
53. Versola-Russo J. Cultural and Demographic Factors of Schizophrenia. *Int J Psychosoc Rehab* 2006;10:89-103.
54. Lindamer LA, Lohr JB, Harris MJ, Jeste DV. Gender, estrogen, and schizophrenia. *Psychopharmacol Bull* 1997;33:221-8.
55. Fearon P, Kirkbride JB, Morgan C, Dazzan P, Morgan K, Lloyd T, et al. Incidence of schizophrenia and other psychoses in ethnic minority groups: results from the MRC AESOP study. *Psychol Med* 2006;36:1541-50.
56. Cantor-Graae E, Selten J-P. Schizophrenia and Migration: A Meta-Analysis and Review. *Am J Psychiatry* 2005;162:12-24.
57. Veling W, Selten J-P, Susser E, Laan W, Mackenbach JP, Hoek HW. Discrimination and the incidence of psychotic disorders among ethnic minorities in The Netherlands. *Int J Epidemiol* 2007;36:761-8.
58. Karlsen S, Nazroo J. The relationship between racism, social class and physical and mental health among different ethnic groups in England. *Ethn Health* 2004;9:46-7.

59. Agerbo E, Byrne M, Eaton WW, Mortenson PB. Marital and Labor Market Status in Long run Schizophrenia. Arch Gen Psychiatry 2004;61:28-33.
60. Marwaha S, Johnson S, Bebbington P. Rates and correlates of employment in people with schizophrenia in the UK, France and Germany. Br J Psychiatry 2007;191:30-7.
61. Usman S, Yousaf A, Asif A, Chaudry HR. Frequency of various Psychiatric disorders managed in indoor setting of a Teaching Hospital. J Fatima Jinnah Med Coll Lahore 2007;1:7-10.
62. Weiser M, van Os J, Reichenferg A, Rabinowitz J, Nahon D, Kravitz E, et al. Social and cognitive functioning, urbanicity and risk for schizophrenia. Br J Psychiatry 2007;191:320-4.
63. van Os J. Does the urban environment cause psychosis? Br J Psychiatry 2004; 184, 287 –8.
64. Cooper B. Immigration and schizophrenia: the social causation hypothesis revisited. Br J Psychiatry 2005; 186: 361-3.
65. Allardyce J, Gilmour H, Atkinson J, Rapson T, Bishop J, McCreadie G. Social fragmentation, deprivation and urbanicity: relation to first-admission rates for psychoses. Br J Psychiatry 2005;187:401-6.

66. Wiles NJ, Zamit S, Bebbington P, Singleton N, Meltzer h, Lewis G. Self-reported psychotic symptoms in the general population. *Br J Psychiatry* 2006;188:519-26.
67. Ruhrmann S, Bechdolf A, Kai-Uwe K, Wagner M, Schultze-Lutter F, Janssen B, et al. Acute effects of treatment for prodromal symptoms for people putatively in late initial prodromal state of psychosis. *Br J Psychiatry* 2007;191:88-95.
68. Stotz-Ingenlath G. Epistemological aspects of Eugen Bleuler's conception of schizophrenia in 1911. *Med Health Care Philos* 2000;3:153-9.
69. Nordgaard J, Arnfred SM, Handest P, Parnas J. The Diagnostic Status of First-Rank Symptoms. *Schizophr Bull* 2008;34:137-54.
70. Bertelsen A. Schizophrenia and Related Disorders: Experience with Current Diagnostic Systems. *Psychopathol* 2002;35:89-93.
71. Breier A, Berg PH. The psychosis of schizophrenia: prevalence, response to atypical antipsychotics, and prediction of outcome. *Biol Psychiatry* 1999;46:361-4.

72. Appelbaum PS, Robbins PC, Roth LH. Dimensional approach to delusions: comparison across types and diagnoses. *Am J Psychiatry* 1999;156:1938-43.
73. Kitamura T, Okazaki Y, Fujinawa A, Takayanagi I, Kasahara Y. Dimensions of schizophrenic positive symptoms: an exploratory factor analysis investigation. *Eur Arch Psychiatry Clin Neurosci* 1998;248:130-5.
74. Velligan DI, Alphas LD. Negative Symptoms in Schizophrenia: The Importance of Identification and Treatment. *Psychiatr Times* 2008;25;3:1-2.
75. Szymanski SR, Cannon TD, Gallacher F, Erwin RJ, Gur RE. Course of treatment response in first-episode and chronic schizophrenia. *Am J Psychiatry* 1996; 153:519-25.
76. Andreasen NC. Negative symptoms in schizophrenia. Definition and reliability. *Arch Gen Psychiatry* 1982; 39:784-8.
77. Hafner H, Maurer K. Are there two types of schizophrenia? True onset and sequence of positive and negative symptoms prior to first admission. In: Marneros A, Andreason NC, Tsuang MT, editors. *Negative Versus Positive Schizophrenia*. Berlin: Springer-Verlag;1991. p.134.

78. McGlashan TH, Fenton WS. The positive-negative distinction in schizophrenia. Review of natural history validators. *Arch Gen Psychiatry* 1992;49:63-72.
79. Ho BC, Nopoulos P, Flaum M, Arndt S, Andreasen NC. Two-year outcome in first-episode schizophrenia: predictive value of symptoms for quality of life. *Am J Psychiatry* 1998;155:1196-201.
80. Cohen AS, Docherty NM. Affective reactivity of speech and emotional experience in patients with schizophrenia. *Schizophr Res* 2004; 69:7-14.
81. Kohler CG, Bilker W, Hagendoorn M, Gur RE, Gur RC. Emotion recognition deficit in schizophrenia: association with symptomatology and cognition. *Biol Psychiatry* 2000;48:127-36.
82. Bell M, Bryson G, Lysaker P. Positive and negative affect recognition in schizophrenia: a comparison with substance abuse and normal control subjects. *Psychiatry Res* 1997;73:73-82.
83. Socco KA, Termine A, Seyal A, Dudas MM, Vessicchio JC, Krishnan-Sarin S et al. Effects of Cigarette Smoking on Spatial Working memory and Attentional Deficits in Schizophrenia. *Arch Gen Psychiatry* 2005;62:649-59.

84. Saykin AJ, Gur RC, Gur RE, Mozley PD, Mozley LH, Resnick SM, et al. Neuropsychological function in schizophrenia. Selective impairment in memory and learning. Arch Gen Psychiatry 1991; 48:618-24.
85. Russell AJ, Munro JC, Jones PB, Hemsley DR, Murray RM. Schizophrenia and the myth of intellectual decline. Am J Psychiatry 1997;154:635-9.
86. Seidman LJ, Buka SL, Goldstein JM, Tsuang MT. Intellectual decline in schizophrenia: evidence from a prospective birth cohort 28 year follow-up study. J Clin Exp Neuropsychol 2006; 28:225-42.
87. Woodberry KA, Giuliano AJ, Seidman LJ. Premorbid IQ in schizophrenia: a meta-analytic review. Am J Psychiatry 2008; 165:579-87.
88. Aylward E, Walker E, Bettes B. Intelligence in schizophrenia: meta-analysis of the research. Schizophr Bull 1984;10:430-59.
89. Hyde TM, Goldberg TE, Egan MF, Lener MC, Weinberger DR. Frontal release signs and cognition in people with schizophrenia, their siblings and healthy controls. Br J Psychiatry 2007; 191: 120-5.

90. Breier A, Schreiber JL, Dyer J, Pickar D. National Institute of Mental Health longitudinal study of chronic schizophrenia. Prognosis and predictors of outcome. Arch Gen Psychiatry 1991;48:239-46.
91. Edwards J, Jackson HJ, Pattison PE. Emotion recognition via facial expression and affective prosody in schizophrenia: a methodological review. Clin Psychol Rev 2002;22:789-832.
92. Schultze-Lutter F, Ruhrmann S, Picker H, Von Reventlow HG, Brockhaus-Dumke A, Klosterkötter J. Basic symptoms in early psychotic and depressive disorders. Br J Psychiatry 2007;191:31-7.
93. Hirsch SR, Jolley A, Barnes T. Are Depressive Symptoms Part of the Schizophrenic Syndrome? In: DeLisi L, editor. Depression in Schizophrenia. Washington DC: American Psychiatric Press; 1990. p.25.
94. Qin P, Nordentoft M. Suicide risk in relation to psychiatric hospitalization: evidence based on longitudinal registers. Arch Gen Psychiatry 2005; 62:427-32.
95. Martin RL, Cloninger CR, Guze SB, Clayton PJ. Frequency and differential diagnosis of depressive syndromes in schizophrenia. J Clin Psychiatry 1985; 46:9-13.

96. Uhlhaas PJ, Phillips WA, Mitchell G, Silverstein SM. Perceptual grouping in disorganized schizophrenia. *Psychiatry Res* 2006;2:105-17.
97. Jakobsen KD, Frederiksen JN, Hansen T, Jansson LB, Parnas J, Werge T. Reliability of clinical ICD-10 schizophrenia diagnoses. *Nord J Psychiatry* 2005;59:209-12.
98. World Health Organization. The ICD-10 classification of mental and behavioural disorders. Geneva: World Health Organization; 1992.
99. Kendler KS, McGuire M, Gruenberg AM, Walsh D. Outcome and family study of the subtypes of schizophrenia in the west of Ireland. *Am J Psychiatry* 1994;151:849-56.
100. Fenton WS, McGlashan TH, Victor BJ, Blyler CR. Symptoms, subtype, and suicidality in patients with schizophrenia spectrum disorders. *Am J Psychiatry* 1997; 154:199-204.
101. Bourgeois M, Swendsen J, Young F, Amador X, Pini S, Cassano GB, et al. Awareness of disorder and suicide risk in the treatment of schizophrenia: results of the international suicide prevention trial. *Am J Psychiatry* 2004; 161:1494-6.
102. Crumlish N, Whitty P, Kamali M, Clarke M, Browne S, Mctigue O, et al. Early insight predicts depression and

- attempted suicide after 4 years in first-episode schizophrenia and schizophreniform disorder. *Acta Psychiatr Scand* 2005; 112:449-55.
103. Deister A, Marneros A. Prognostic value of initial subtype in schizophrenic disorders. *Schizophr Res* 1994; 12:145-57.
 104. Taylor MA, Abrams R. Catatonia. Prevalence and importance in the manic phase of manic-depressive illness. *Arch Gen Psychiatry* 1977; 34:1223-25.
 105. Peralta V, Cuesta MJ, Serrano JF, Mata I. The Kahlbaum syndrome: a study of its clinical validity, nosological status, and relationship with schizophrenia and mood disorder. *Compr Psychiatry* 1997; 38:61-7.
 106. Fenton WS, McGlashan TH. Natural history of schizophrenia subtypes. I. Longitudinal study of paranoid, hebephrenic, and undifferentiated schizophrenia. *Arch Gen Psychiatry* 1991; 48:969-77.
 107. Isohanni M, Lauronen E, Moilanen K, Isohanni I, Kemppainen L, Koponen H, et al. Predictors of schizophrenia. Evidence from Northern Finland 1966 Birth Cohort and other sources. *Br J Psychiatry* 2005;87:4-7.

108. Davies EJ. Developmental aspects of schizophrenia and related disorders: possible implications for treatment strategies. *Adv Psychiatr Treat* 2007; 13: 384-91.
109. Tsuang M. Schizophrenia: genes and environment. *Biol Psychiatry* 2000;47:210-20.
110. Stefansson H, Sigurdsson E, Steinthorsdottir V, Bjornsdottir S, Sigmundsson T, Ghosh S, et al. Neuregulin 1 and susceptibility to schizophrenia. *Am J Hum Genet* 2002; 71:877-92.
111. Stefansson H, Sarginson J, Kong A, Yates P, Steinthorsdottir V, Gudfinnsson E, et al. Association of neuregulin 1 with schizophrenia confirmed in a Scottish population. *Am J Hum Genet* 2003; 72:83-7.
112. Sanders AR, Duan J, Levinson DF, Shi J, He D, Hou C, et al. No significant association of 14 candidate genes with schizophrenia in a large European ancestry sample: implications for psychiatric genetics. *Am J Psychiatry* 2008; 165:497-506.
113. O'Donovan MC, Craddock N, Norton N, Williams H, Pierce T, Moskvina V, et al. Identification of loci associated with

schizophrenia by genome-wide association and follow-up.
Nat Genet 2008;40:1053-5.

114. Walsh T, McClellan JM, McCarthy SE, Addington AM, Pierce SB, Cooper GM, et al. Rare structural variants disrupt multiple genes in neurodevelopmental pathways in schizophrenia. Science 2008;320:539-43.
115. Allen NC, Bagade, S McQueen M, Ioannidis JP, Kavvoura FK, Khoury MJ, et al. Systematic meta-analyses and field synopsis of genetic association studies in schizophrenia: the SzGene database. Nat Genet 2008; 40:827-34.
116. Sipos A, Rasmussen F, Harrison G, Tynelius P, Lewis G, Leon DA, et al. Paternal age and schizophrenia: a population based cohort study. BMJ 2004; 329:1070-3.
117. Munk-Jorgensen P, Ewald, H. Epidemiology in neurobiological research: exemplified by the influenza-schizophrenia theory. Br J Psychiatry 2001;40:s30-2.
118. Susser E, Neugebauer R, Hoek HW, Brown AS, Lin S, Labovitz D, et al. Schizophrenia after prenatal famine. Further evidence. Arch Gen Psychiatry 1996;53:25-31.

119. St Clair D, Xu M, Wang P, Yu Y, Fang Y, Zhang F, et al. Rates of adult schizophrenia following prenatal exposure to the Chinese famine of 1959-1961. *JAMA* 2005;294:557-62.
120. Bresnahan M, Schaefer CA, Brown AS, Susser ES. Prenatal determinants of schizophrenia: what we have learned thus far? *Epidemiol Psychiatr Soc* 2005;14:194-7.
121. McNeil TF, Cantor-Graae E, Ismail B. Obstetric complications and congenital malformation in schizophrenia. *Brain Res Brain Res Rev* 2000;31:166-78.
122. Buhler B, Hambrecht M, Loffler W, van der Heiden W, Häfner H. Precipitation and determination of the onset and course of schizophrenia by substance abuse--a retrospective and prospective study of 232 population-based first illness episodes. *Schizophr Res* 2002; 54:243-51.
123. Selton JP, Cantor-Grace E, Kahn RS. Migration and schizophrenia. *Curr Opin Psychiatry* 2007;20:111-5.
124. Davies G, Welham J, Chant D, Torrey EF, McGrath J. A systematic review and meta-analysis of Northern Hemisphere season of birth studies in schizophrenia. *Schizophr Bull* 2003;29:587-93.

125. Brown AS. Prenatal infection as a risk factor for schizophrenia. *Schizophr Bull* 2006;32:200-2.
126. Schenkel LS, Spaulding WD, Dilillo D, Silverstein SM. Histories of childhood maltreatment in schizophrenia: Relationships with premorbid functioning, symptomatology and cognitive deficits. *Schizophr Res* 2005;76:273-86.
127. Bentall RP, Fernyhough C, Morrison AP, Lewis S, Corcoran R. Prospects for cognitive-developmental account of psychotic experiences. *Br J Clin Psychol* 2007;46:155-73.
128. Ferdinand RF, Sondeijker F, van der Ende J, Selton JP, Huizink A, Verhulst FC. Cannabis use predicts future psychotic symptoms, and vice versa. *Addiction* 2005;100:612-8.
129. Seeman P. Targeting the dopamine D2 receptor in schizophrenia. *Expert Opin Ther Targets* 2006;10:515-31.
130. Weinberger DR, Egan MF, Bertolino A, Callicott JH, Mattay VS, Lipska BK, et al. Prefrontal neurons and the genetics of schizophrenia. *Biol Psychiatry* 2001; 50:825-44.
131. Shenton ME, Dickey CC, Frumin M, McCarley RW. A review of MRI findings in schizophrenia. *Schizophr Res* 2001; 49:1-52.

132. McGlashan TH, Hoffman RE. Schizophrenia as a disorder of developmentally reduced synaptic connectivity. Arch Gen Psychiatry 2000; 57:637-48.
133. Hendren RL, De Backer I, Pandina GJ. Review of neuroimaging studies of child and adolescent psychiatric disorders from the past 10 years. J Am Acad Child Adolesc Psychiatry 2000; 39:815-28.
134. Boos HB, Cahn W, Appels MC, Sitskoorn MM, Hulshoff Pol HE, Schnack HG, et al. Brain volumes in parents of patients with schizophrenia. Schizophr Bull 2005;31:382-90.
135. van Haren NE, Picchioni MM, McDonald C, Marshall N, Davis N, Ribchester T, et al. A controlled study of brain structure in monozygotic twins concordant and discordant for schizophrenia. Biol Psychiatry. 2004;56:454-61.
136. Liberman JA. Neurobiology and the natural history of schizophrenia. J Clin Psychiatry 2006 ;67:14-8.
137. Alem A, Kebede D, Fekadu A, Shibre T, Fekadu D, Beyero T, et al. Course and Outcome of Schizophrenia in a Predominantly Treatment-Naive Cohort in Rural Ethiopia. Schizophr Bull 2009; 35:646-54.

138. Robinson DG, Woerner MG, McMeniman M, Mendelowitz A, Bilder RM. Symptomatic and functional recovery from a first episode of schizophrenia or schizoaffective disorder. *Am J Psychiatry* 2004; 161:473–9.
139. Hellidín L, Kane JM, Karilampi U, Norlander T, Archer T. Remission in prognosis of functional outcome: a new dimension in the treatment of patients with psychotic disorders. *Schizophr Res* 2007;93:160-8.
140. Rabinowitz J, Harvey PD, Eerdeken M, Davidson M. Premorbid functioning and treatment response in recent-onset schizophrenia. *Br J Psychiatry* 2006;189:31-5.
141. Schimmelmann BG, Huber CG, Lambert M, Cotton S, McGorry PD, Conus P. Impact of duration of untreated psychosis on pre-treatment, baseline, and outcome characteristics in an epidemiological first-episode psychosis cohort. *J Psychiatr Res* 2008;42:982-90.
142. Newman SC, Bland RC. Mortality in a cohort of patients with schizophrenia: a record linkage study. *Can J Psychiatry* 1991; 36:239-45.

143. Marder SR, Essock SM, Miller AL, Buchanan RW, Casey DE, Davis JM, et al. Physical health monitoring of patients with schizophrenia. *Am J Psychiatry* 2004; 161:1334-49.
144. Ponnudurai R, Jayakar J, Sathya-Sekaran BWC. Assessment of mortality and marital status of schizophrenic patients over a period of 13 years. *Ind J Psychiatry* 2006;48:84-7.
145. Palmer BA, Pankratz VS, Bostwick JM. The Lifetime Risk of Suicide in Schizophrenia: A Reexamination. *Arch Gen Psychiatry* 2005;62:247-53.
146. Almond S, Knapp M, Francois C, Toume M, Brugha T. Relapse in schizophrenia: costs, clinical outcomes and quality of life. *Br J Psychiatry* 2004; 184: 346-51.
147. McPhillips MA, Kelly FJ, Barnes TR, Duke PJ, Gene-Cos N, Clark K. Detecting comorbid substance misuse among people with schizophrenia in the community: a study comparing the results of questionnaires with analysis of hair and urine. *Schizophr Res* 1997; 25:141-8.
148. de Leon J, Tracy J, McCann E, McGrory A, Diaz FJ. Schizophrenia and tobacco smoking: a replication study in

- another US psychiatric hospital. *Schizophr Res* 2002;56:55-65.
149. Carney CP, Jones L, Woolson RF. Medical comorbidity in women and men with schizophrenia: a population-based controlled study. *J Gen Intern Med* 2006; 21:1133-7.
 150. Rehman I, Farooq S. Cannabis abuse in patients with Schizophrenia: pattern and effects on symptomatology. *J Coll Physician Surg Pak* 2007;17:158-61.
 151. Chwastiak LA, Rosnheck RA, McEvoy JP, Keefe RS, Swartz MS, Liberman JA. Interrelationship of Psychiatric Symptom Severity, Medical Comorbidity, and Functioning in Schizophrenia. *Psychiatr Serv* 2006;57:1102-9.
 152. Lenroot R, Bustillo JR, Lauriello J, Keith SJ. Integration of Care: Integrated Treatment of Schizophrenia. *Psychiatr Serv* 2003;54:1499-1507.
 153. Nasrallah HA, Targum SD, Tandon R, McCombs JS, Ross R. Defining and Measuring Clinical Effectiveness in the Treatment of Schizophrenia. *Psychiatr Serv* 2005; 56:273-82.
 154. Gaebel W, Weinmann S, Sartorius N, Rutz W, McIntyre JS. Schizophrenia practice guidelines: international survey and comparison. *Br J Psychiatry* 2007;187:248-55.

155. Royal Australian and New Zealand College of Psychiatrists Clinical Practice Guidelines Team for the Treatment of Schizophrenia and Related Disorders. Aust NZ J Psychiatry 2005;39:1-30.
156. Lewis SW, Tarrier N, Drake RJ. Integrating non-drug treatments in early schizophrenia. Br J Psychiatry 2005; 187: 65-71.
157. Marder SR, Wirshing WC, Mintz J, McKenzie J, Johnston K, Eckman TA, et al. Two-year outcome of social skills training and group psychotherapy for outpatients with schizophrenia. Am J Psychiatry 1996; 153:1585-92.
158. Habib N, Khan SD, Kingdon D, Naeem F. Preliminary evaluation of culturally adapted CBT for psychosis. imPACT 2012;3:7.
159. Lauriello J, Bustillo J, Keith SJ. A critical review of research on psychosocial treatment of schizophrenia. Biol Psychiatry 1999; 46:1409-17.
160. Miyamoto S, Duncan GE, Marx CE, Liberman JA. Treatment of schizophrenia : a critical review of pharmacology and mechanism of action of antipsychotic drugs. Mol Psychiatry 2005;10:79-104.

161. Naseem R, Chaudry HR, Keshavan MS. An Update on management of Schizophrenia. Med Channel 2001;7:15-8.
162. Lieberman JA, Scott Stroup T, McEvoy JP, Swar MS, Rosenheck RA, Perkins DO. Effectiveness of Antipsychotic Drugs in Patients with Chronic Schizophrenia. N Engl J Med 2005; 353:1209-23.
163. Stargardt T, Weinbrenner S, Busse R, Juckel G, Gericke CA. Effectiveness and cost of atypical versus typical antipsychotic treatment for schizophrenia in routine care. J Ment Health Policy Econ 2008; 11: 89-97.
164. Taylor DM, Douglas-Hall P, Olofinjana B, Whiskey E, Thomas A. Reasons for discontinuing clozapine: matched, case–control comparison with risperidone long-acting injection. Br J Psychiatry 2009;194:165-7.
165. Homel P, Casey D, Allison DB. Changes in body mass index for individuals with and without schizophrenia, 1987-1996. Schizophr Res 2002; 55:277-84.
166. McEvoy JP, Meyer JM, Goff DC, Nasrallah HA, Davis SM, Sullivan L, et al. Prevalence of the metabolic syndrome in patients with schizophrenia: baseline results from the Clinical Antipsychotic Trials of Intervention Effectiveness (CATIE)

- schizophrenia trial and comparison with national estimates from NHANES III. *Schizophr Res* 2005; 80:19-32.
167. Cohen D, Stolk RP, Grobbee DE, Gispen-de Wied CC. Hyperglycemia and diabetes in patients with schizophrenia or schizoaffective disorders. *Diabetes Care* 2006; 29:786-91.
 168. Mackin P, Watkinson HM, Young AH. Prevalence of obesity, glucose homeostasis disorders and metabolic syndrome in psychiatric patients taking typical or atypical antipsychotic drugs: a cross-sectional study. *Diabetologia* 2005; 48:215-21.
 169. Newcomer JW, Hennekens CH. Severe mental illness and risk of cardiovascular disease. *JAMA* 2007; 298:1794-96.
 170. Goff DC, Sullivan LM, McEvoy JP, Meyer JM, Nasrallah HA, Daumit GL, et al. A comparison of ten-year cardiac risk estimates in schizophrenia patients from the CATIE study and matched controls. *Schizophr Res* 2005; 80:45-53.
 171. Glassman AH. Schizophrenia, antipsychotic drugs, and cardiovascular disease. *J Clin Psychiatry* 2005;66:5-10.
 172. US Food and Drug Administration. Information for Healthcare Professionals: Haloperidol. [Online]. 2007. [cited 2009 November 26]. Available from URL:<http://www.fda.gov/cder/drug/InfoSheets/HCP/haloperidol.htm>.

173. Ali W, Maqsood N, Rehman W. Schizophrenia and drug non-compliance. *Professional Med J* 2006;13:423-30.
174. Taj R, Khan S. A study of reasons of non-compliance to psychiatric treatment. *J Ayub Med Coll Abbottabad* 2005;17:26-8.
175. Copolov DL. Psychoses: a primary care perspective. *Med J Aust* 1998;168:129–35.
176. Tooth GC, Brooke M. Trends in the mental hospital population and their effect on future planning. *Lancet* 1961;1:710-3.
177. Johnstone EC, Owens DGC, Gold A, Crow TJ, Macmillan JF. Schizophrenic patients discharged from hospital-A follow-up study. *Br J Psychiatry* 1984;145:586-90.
178. Burns T, Kendrick T. The primary care of patients with schizophrenia: a search for good practice. *Br J Gen Pract* 1997;47:515–20.
179. Carr VJ. The role of the general practitioner in the treatment of schizophrenia: general principles. *Med J Aust* 1997;166:91–4.
180. Keks NA, Altson BM, Sacks TL, Hustig HH, Tanaghow A. Collaboration between general practice and community

- psychiatric services for people with chronic mental illness.
Med J Aust 1997;167:266–71.
181. Copolov DL. Psychoses: a primary care perspective. Med J Aust 1998;168:129–35.
 182. Falloon IRH, Fadden G. Integrated mental health care. Cambridge: Cambridge University Press;1993.
 183. King M, Nazareth I. Community care of patients with schizophrenia: the role of the primary health care team. Br J Gen Pract 1996;46:231–7.
 184. Nazareth ID, King MB. Schizophrenia: community care and the family physician. Int Rev Psychiatry 1992;4:267–72.
 185. Jorm AF. Mental health literacy: public knowledge and beliefs about mental disorders. Br J Psychiatry 2000;177:396–401.
 186. Carr VJ, Lewin TJ, Barnard RE, Walton JM, Allen JL, Constable PM, et al. Attitudes and roles of general practitioners in the treatment of schizophrenia compared with community mental health staff and patients. Soc Psychiatry Psychiatr Epidemiol 2004;39:78–84.
 187. Collins KA, Wolfe VV, Fisman S, DePace J, Steele M. Managing Depression in primary care. Can Fam Physician 2006;52:878-9.

188. Barbato A. World Health Organization: Schizophrenia and public health. [Online] 1998 [Cited on January 15, 2012]. Available from URL: http://www.who.int/mental_health/media/en/55.pdf
189. Valenstein M, Klinkman M, Becker S, Blow FC, Barry KL, Sallar A, et al. Concurrent treatment of patients with depression in the community: provider practices, attitudes, and barriers to collaboration. *J Fam Pract* 1999;48:180-7.
190. Katon W, VonKorff M, Lin E, Walker E, Simon GE, Bush T, et al. Collaborative management to achieve treatment guidelines: impact on depression in primary care. *JAMA* 1995;273:1026-31.
191. Parkes CM, Brown GW, Monck EM. The general practitioner and the schizophrenic patient. *BMJ* 1962;i:972-6.
192. King M, Nazareth I. Community care of patients with schizophrenia: the role of the primary health care team. *Br J Gen Pract* 1996;46:231–7
193. Wang PS, Berglund P, Kessler RC. Recent care of common mental disorders in the United States: prevalence and conformance with evidenced-based recommendations. *J Gen Intern Med* 2000;15:284-92.

194. Simon A, Umbricht D, IGPS Group. Early psychosis in primary care: implications of the international GP study (IGPS) on service development. *Schizophr Res* 2004;70:38-9.
195. Lewin TJ, Carr VJ. Rates of treatment of schizophrenia by general practitioners. A pilot study. *Med J Aust* 1998;168:166-9.
196. Naqvi HA, Sabzwari S, Hussain S, Islam M, Zaman M. General practitioners' awareness and management of common psychiatric disorders: a community-based survey from Karachi, Pakistan. *East Mediterr Health J* 2012;18:446-53.
197. Verdoux H, Cougnard A, Grolleau S, Besson R, Delcroix F. A survey of general practitioners' knowledge of symptoms and epidemiology of schizophrenia. *Eur Psychiatry* 2006;21:238-44.
198. Practice guidelines for the treatment of patients with schizophrenia. American Psychiatric Association. *Am J Psychiatry* 1997;154:1-63.
199. PennDL, Corrigan PW. The effects of stereotype suppression on psychiatric stigma. *Schizophr Res* 2002;55:269-76.
200. Mc Gorry PD, McKenzie D, Jackson HJ. Can we improve the diagnostic efficiency, and predictive power of prodromal symptoms for schizophrenia? *Schizophr Res* 2000;42:91-100.

201. Oud MJT, Schuling J, Slooff CJ, Groenier KH, Dekker JH, de Jong BM. Care for patients with severe mental illness: the general practitioner's role perspective. *BMC Family Practice* 2009, 10:29-36.
202. Report of the Subcommittee on Mental Health and Substance Abuse. Ninth Five- year Plan (1998–2003). Prospective plan 2003–2013. Islamabad, Planning Commission, Government of Pakistan;1998.
203. Irfan M. The Concept of Mental Health Policy and its Journey from Development to implementation in Pakistan. *KUST Med J* 2010; 2(2): 64-8.
204. Norton JL, Rivoiron-Besset E, David M, Jaussent I, Prudhomme C, Boulenger JP, et al. Role of the General Practitioner in the care of patients recently discharged from the hospital after a first psychotic episode: Influence of length of stay. *Prim Care Companion CNS Disord* 2011; 13(5): PCC.11m01180. doi: 10.4088/PCC.11m01180.
205. Capdevielle D, Ritchie K. The long and the short of it: are shorter periods of hospitalisation beneficial? *Br J Psychiatry* 2008;192(3):164–5.

206. Dietrich AJ, Oxman TE, Williams JW Jr, et al. Re-engineering systems for the treatment of depression in primary care: cluster randomised controlled trial. *BMJ* 2004;329(7466):602.
207. Gilbody S, Whitty P, Grimshaw J, et al. Educational and organizational interventions to improve the management of depression in primary care: a systematic review. *JAMA* 2003;289(23):3145–51.
208. Upshur C, Weinreb L. A survey of primary care provider attitudes and behaviors regarding treatment of adult depression: what changes after a collaborative care intervention? *Prim Care Companion J Clin Psychiatry*. 2008;10(3):182–6.
209. Beck AT, Steer RA. Manual for the revised Beck Depression Inventory. San Antonio, Tex: Psychological Corporation; 1993.
210. Hamilton M. Development of a rating scale for primary depressive illness. *J Neurol Neurosurg Psychiatry* 1967;2:56-62.

CHAPTER 5: APPENDIX 'A'

KNOWLEDGE AND PRACTICES OF GENERAL PRACTITIONERS OF DISTRICT PESHAWAR ABOUT SCHIZOPHRENIA

5.1 PROFORMA

Direction: *Kindly fill the proforma and where required put a ✓ in the appropriate box*

Serial _____ Number: _____ Contact _____ Details: _____

❖ GENERAL INFORMATION

- Name: _____
- Gender:
 - Male ☐
 - Female ☐
- How many patients with an established diagnosis of schizophrenia do you treat annually?
 - None ☐
 - 1–2 ☐
 - 3–5 ☐
 - 6–9 ☐
 - More than 10 ☐
- How much time do you generally take for a consultation by someone with schizophrenia?
 - Less than 10 min ☐
 - 10–20 min ☐
 - 20–30 min ☐
 - More than 30 min ☐

- How many patients in whom you suspect the onset of schizophrenia do you see in your practice?
- None ☐
 - 1–2/year ☐
 - 3–5/year ☐
 - More than 5/year ☐

❖ **KNOWLEDGE**

- Which of the following do you think are the most frequent symptoms of Schizophrenia? (more than one answer allowed)
- Hallucinations/delusions ☐
 - Social withdrawal ☐
 - Psychosomatic complaints ☐
 - Suicidality ☐
 - Depression/anxiety ☐
 - Bizarre behaviour ☐
 - Drug misuse ☐
 - Conflicts with parents/teachers/employers ☐
 - Functional decline (school/work) ☐
- Do you think that a first episode of schizophrenia is preceded by early warning signs?
- Yes ☐
 - No ☐
- What therapy in your thinking is ideal for a patient with a suspected first schizophrenic episode (independent of whether you treat these patients yourself)? (more than one answer allowed)
- Psychotherapy ☐
 - Pharmacotherapy ☐
 - Family therapy ☐
 - Observe and wait only ☐

➤ For how long should antipsychotic medication be maintained after a first schizophrenic episode?

- Few days ☐
- 3–4 weeks ☐
- 1–6 months ☐
- 6–12 months ☐
- 12–24 months ☐
- 3–5 years ☐

➤ For how long should antipsychotic medication be maintained in patients with multiple episodes of illness after the remission of an episode?

- Few days ☐
- 3–4 weeks ☐
- 1–6 months ☐
- 6–12 months ☐
- 12–24 months ☐
- At least 3–5 years ☐

➤ How high do you estimate the relapse risk of untreated patients during the first year after a first schizophrenic episode?

_____ %

➤ Which are the two clinically most relevant side-effects of antipsychotic treatment?

❖ PRACTICES

- What do you generally do to confirm the diagnosis? (more than one answer allowed)
 - Personal history ☐
 - Family history ☐
 - Information from significant others(teacher/employer) ☐
 - Observation over several days and weeks ☐
 - Observation over several months ☐
 - Neurological assessment ☐
 - Neuropsychological assessment ☐
 - Other examinations(radiographic,electrophysiological) ☐
 - Laboratory tests ☐
 - Urine testing for drug abuse ☐
 - Consultation with/referral to a specialist ☐
 - Other (specify) _____

- Are these patients treated by you alone, or in collaboration with other specialists or institutions? (more than one answer allowed)
 - Treatment exclusively in my clinic ☐
 - Occasional/regular consultation with a specialist to reassess/advise ☐
 - Referral to a specialist for initial diagnosis and to establish the medication regimen, continuation of treatment in my clinic ☐
 - Referral to a specialist/psychiatric out-patient department and complete handover for treatment ☐

- What medications are commonly used in patients with schizophrenia and what are the doses?

_____ (name) _____ mg/day

_____ (name) _____ mg/day

_____ (name) _____ mg/day

_____ (name) _____ mg/day

- Based on your experience, how do you judge the prognosis of a treated patient after a first schizophrenic episode? (more than one answer allowed)

- The prognosis may be favourable; one single episode with maintenance of performance level is possible ☐
- Mostly several episodes with possible maintenance of performance level ☐
- Mostly several episodes with progressive decline of performance level and severe course of illness ☐

- Did you participate in continuing education on schizophrenia or early schizophrenic psychosis in the past few months?

- No ☐
- Yes; indicate name, place and date of education ☐

Thank you for your participation